

University of Groningen

Do BL Lac objects fit on the fundamental plane?

Plotkin, Richard; Markoff, Sera; Kelley, Brandon; Koerding, Elmar; Trager, Scott

Published in:

38th COSPAR Scientific Assembly. Held 18-15 July 2010, in Bremen, Germany, p.6

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Plotkin, R., Markoff, S., Kelley, B., Koerding, E., & Trager, S. (2010). Do BL Lac objects fit on the fundamental plane? In *38th COSPAR Scientific Assembly. Held 18-15 July 2010, in Bremen, Germany, p.6* (pp. 6) <http://adsabs.harvard.edu/abs/2010cosp...38.2459P>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Research in Astrophysics from Space (E)

Accretion on Compact Objects and Fast Phenomena in Multiwavelength Era (E13)

DO BL LAC OBJECTS FIT ON THE FUNDAMENTAL PLANE?

Richard Plotkin, r.m.plotkin@uva.nl

University of Amsterdam, Amsterdam, The Netherlands

Sera Markoff, s.b.markoff@uva.nl

University of Amsterdam, Amsterdam, The Netherlands

Brandon Kelley, bckelly@cfa.harvard.edu

Harvard-Smithsonian Center for Astrophysics, United States

Elmar Koerding, elmar@koerding.eu

AIM - Universite Paris 7 CEA Saclay, Gif-sur-Yvette, France

Scott Trager, sctrager@astro.rug.nl

Kapteyn Astronomical Institute, University of Groningen, The Netherlands

The fundamental plane of black hole accretion—a correlation between radio luminosity, X-ray luminosity, and black hole mass—expresses the connections between accretion physics over at least eight orders of magnitude in black hole mass. We first present an updated regression analysis of the fundamental plane utilizing a Bayesian approach. We then present new results on our attempts to place BL Lac objects on the fundamental plane. BL Lac objects are interpreted in the standard model as low-luminosity radio galaxies viewed nearly along the axis of a relativistic jet. They might be excellent high-mass analogs ($\sim 10^8$ - 10^9 solar masses) to hard-state X-ray binaries because their spectral energy distributions are jet dominated. Also, they likely have weak broad emission line regions, meaning their environments are relatively uncomplicated compared to other classes of Active Galactic Nuclei (AGN) with similarly massive black holes. We consider around 10^2 BL Lac objects here (taken from a much larger 723 object SDSS BL Lac sample) for which we measure black hole masses, and for which we have radio, optical, and X-ray flux measurements. Even just this subset of 10^2 BL Lac objects is larger than entire XRB/AGN samples used in previous fundamental plane studies. Our BL Lacs generally fit on the fundamental plane, albeit with scatter. We investigate potential sources of this scatter, such as the importance of relativistic beaming. We also discuss if objects deviate from the plane as a function of their synchrotron cutoff frequency, which could be interpreted as variations in the strengths of their broad emission line regions (i.e., environment) and/or in accretion rate.